

**NTP Technical Report
on Toxicity Studies of**

Glyphosate
(CAS No. 1071-83-6)

**Administered in Dosed Feed
to F344/N Rats and B6C3F₁ Mice**

**Po C. Chan, PhD, and Joel F. Mahler, DVM,
Study Scientists
National Toxicology Program
Post Office Box 12233
Research Triangle Park, NC 27709**

**NIH Publication 92-3135
July 1992**

**United States Department of Health and Human Services
Public Health Service
National Institutes of Health**

CONTRIBUTORS

The NTP report on the toxicity studies of glyphosate is based on disposition studies conducted at the College of Pharmacy, University of Arizona, Tucson, AZ, in November, 1987; 13-week studies performed between May and September, 1988, at Southern Research Institute, Birmingham, AL; and 14-day studies performed in 1990 at the National Institute of Environmental Health Sciences, Research Triangle Park, NC.

National Toxicology Program

*Evaluated experiment, interpreted results,
and reported findings*

Po C. Chan, PhD
Joel F. Mahler, DVM
Study Scientists

John R. Bucher, PhD
Leo T. Burka, PhD
Rajendra S. Chhabra, PhD
Michael P. Dieter, PhD
Michael R. Elwell, DVM, PhD
H.B. Matthews, PhD
Morrow B. Thompson, DVM, PhD
Errol Zeiger, PhD

Coordinated report preparation

Jane M. Lambert, BS
Diane Overstreet, BS
Kristine Witt, MS
Oak Ridge Associated Universities

NTP Pathology Review

Evaluated slides and prepared pathology report

Sondra Grumbein, DVM, PhD,
Pathology Associates, Inc.
Michael R. Elwell, DVM, PhD
National Toxicology Program

Southern Research Institute

Principal contributors

J.D. Prejean, PhD
Principal Investigator
H. Giles, DVM
A.G. Manus, DVM,
L.M. Thigpen, DVM
R. Thompson, DVM

University of Arizona, College of Pharmacy

Principal contributors, disposition studies

I. Glenn Sipes, PhD
Christy Duerson, MS

Experimental Pathology Laboratories, Inc.

Provided pathology quality assurance

Jerry F. Hardisty, DVM

Environmental Health Research and Testing, Inc.

*Provided sperm morphology and
reproductive toxicology evaluation*

Dushant K. Gulati, PhD
Teresa Cocanougher, BA
Susan Russell, BA

Analytical Sciences, Inc.

Provided statistical analysis

Steven Seilkop, MS
Janet Teague, MS

CONTENTS

CONTRIBUTORS	2
TABLE OF CONTENTS	3
ABSTRACT	5
PEER REVIEW PANEL	7
SUMMARY OF PEER REVIEW COMMENTS	8
INTRODUCTION	9
MATERIALS AND METHODS	11
Procurement and Characterization of Glyphosate	11
Disposition Studies	11
13-Week Study Design	12
Reproductive Toxicity	13
Study of the Mechanism of Induction of Salivary Gland Lesions by Glyphosate	15
Genetic Toxicity Studies	16
Statistical Methods	16
Quality Assurance	17
RESULTS	18
Disposition Studies	18
13-Week Studies in F344/N Rats	19
13-Week Studies in B6C3F ₁ Mice	23
Mechanism of Induction of Salivary Gland Lesions	28
Genetic toxicology	33
DISCUSSION	34
REFERENCES	37
TABLES	
Table 1	Experimental Design and Materials and Methods in the 13-Week Studies of Glyphosate
	14
Table 2	Treatment Groups in the Study to Determine the Mechanism of Induction of Salivary Gland Lesions by Glyphosate
	15
Table 3	Cumulative Percentage of Oral or I.V. Dose of Glyphosate Eliminated in Urine and Feces
	18
Table 4	Percentage of Dose in Tissues Following Oral Administration of Glyphosate at 5.6 mg/kg
	18

Table 5	Survival, Weight Gain, and Feed Consumption of F344/N Rats in the 13-Week Dosed Feed Study of Glyphosate	19
Table 6	Incidence and Severity of Cytoplasmic Alteration of the Parotid and Submandibular Salivary Glands (Combined) in F344/N Rats in the 13-Week Dosed Feed Study of Glyphosate	22
Table 7	Survival, Weight Gain, and Feed Consumption of B6C3F ₁ Mice in the 13-Week Dosed Feed Study of Glyphosate	23
Table 8	Incidence and Severity of Cytoplasmic Alteration of the Parotid Salivary Gland in B6C3F ₁ Mice in the 13-Week Dosed Feed Study of Glyphosate	23
Table 9	Feed Consumption and Weight Gain of F344/N Rats in the 14-Day Mechanism Study of Glyphosate	28
Table 10	Salivary Gland Weights of F344/N Rats in the 14-Day Mechanism Study of Glyphosate	29
Table 11	Incidence and Severity of Cytoplasmic Alteration of the Salivary Glands of F344/N Rats in the 14-Day Mechanism Study of Glyphosate	29

FIGURES

Figure 1	Blood Levels of ¹⁴ C-Glyphosate Following Oral Administration of ¹⁴ C-Glyphosate at 5.6 or 56 mg/kg	20
Figure 2	Level of Radioactivity in Blood after a Single I.V. Dose of 5.6 mg/kg Glyphosate	20
Figure 3	Body Weights of F344/N Rats Exposed to Glyphosate by Dosed Feeding for 13 Weeks	21
Figure 4	Body Weights of B6C3F ₁ Mice Exposed to Glyphosate by Dosed Feeding for 13 Weeks	27


PLATES

Plates 1-2	24
Plates 3-5	30

APPENDICES

Appendix A	Organ Weights and Organ-to-Body-Weight Ratios	A-1
Appendix B	Hematology and Clinical Chemistry Results	B-1
Appendix C	Reproductive Tissue Evaluations and Estrous Cycle Length	C-1
Appendix D	Genetic Toxicology	D-1

Glyphosate



Molecular Formula: C₃H₈NO₅P

CAS Number: 1071-83-6

Molecular Weight: 169.1

Synonyms: Glyphosate, technical grade; Glycine, N-(phosphonomethyl); N-phosphonomethyl glycine; N-(phosphonomethyl)glycine; MON 0573; MON 2139.

ABSTRACT

Glyphosate is a systemic, broad-spectrum, post-emergence herbicide used for non-selective weed control. It was selected for study because of its widespread use, potential for human exposure, and the lack of published reports concerning comprehensive toxicity or carcinogenicity evaluations.

Chemical disposition, 13-week toxicity, and mutagenicity studies of glyphosate were conducted. In disposition studies, male F344/N rats were administered an oral dose (5.6 or 56 mg/kg) of ¹⁴C-glyphosate. Blood, urine, fecal, and tissue samples were collected and analyzed for radioactivity. Within 72 hours after glyphosate dosing, 20-30% of the administered radioactivity was eliminated via urine, 70-80% via feces, and about 1% of the radioactivity remained in the tissues. Studies following oral, intravenous, and intraperitoneal administration of glyphosate indicated that the urinary radioactivity represented the amount of glyphosate absorbed and that the fecal radioactivity represented the amount unabsorbed from the gastrointestinal tract.

In the 13-week toxicity studies, groups of 10 male and female F344/N rats and B6C3F₁ mice were administered glyphosate in feed at 0, 3125, 6250, 12500, 25000, or 50000 ppm. Glyphosate administration induced increases in serum bile acids, alkaline phosphatase, and alanine aminotransferase activities in rats, suggesting mild toxicity to the hepatobiliary system. Clinical pathology measurements were not performed with mice. No histopathologic lesions were observed in the livers of rats or mice. There was no evidence of adverse effects on the reproductive system of rats or mice. Cytoplasmic alteration was observed in the parotid and submandibular salivary glands of rats and parotid salivary glands in mice. The salivary gland

effects of glyphosate were demonstrated to be mediated through an adrenergic mechanism which could be blocked by the adrenergic antagonist, propranolol.

Glyphosate was not mutagenic in *Salmonella*, and did not induce micronuclei in mice. The no-observed-adverse-effect level (NOAEL) for the salivary gland lesions was 3125 ppm in the diet for mice. A NOAEL could not be determined from the rat study.

PEER REVIEW

Peer Review Panel

The members of the Peer Review Panel who evaluated the draft report on the toxicity studies on glyphosate on July 10, 1991, are listed below. Panel members serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, panel members act to determine that the design and conditions of the NTP studies were appropriate and to ensure that the toxicity study report presents the experimental results and conclusions fully and clearly.

National Toxicology Program's Board of Scientific Counselors Technical Reports Review Subcommittee

Paul T. Bailey, PhD
Mobil Oil Corporation
Toxicology Division
Princeton, NJ

Louis S. Beliczky, MS, MPH
Director of Industrial Hygiene
Department of Industrial Hygiene
United Rubber Workers Intl. Union
87 South High Street
Akron, OH

Gary P. Carlson, PhD
Department of Pharmacology and Toxicology
Purdue University
West Lafayette, IN

Harold Davis, DVM, PhD
School of Aerospace Medicine
Brooks Air Force Base, TX

Robert H. Garman, DVM
Consultants in Veterinary Pathology
Murrysville, PA

Jay I. Goodman, PhD
Department of Pharmacology and Toxicology
Michigan State University
East Lansing, MI

*Unable to attend

David W. Hayden, DVM, PhD
Department of Veterinary Pathobiology
College of Veterinary Medicine
University of Minnesota
St. Paul, MN

Daniel S. Longnecker, MD, Chair
Department of Pathology
Dartmouth Medical School
Hanover, NH

Curtis D. Klaassen, PhD
Department of Pharmacology and Toxicology
University of Kansas Medical Center
Kansas City, KS

Barbara McKnight, PhD
Department of Biostatistics
University of Washington
Seattle, WA

*Ellen K. Silbergeld, PhD
University of Maryland Medical School
Baltimore, MD

Lauren Zeise, PhD
California Department of Health Services
Berkeley, CA

Summary of Peer Review Comments

On July 9 and 10, 1991, the Technical Reports Review Subcommittee of the Board of Scientific Counselors for the National Toxicology Program met in Research Triangle Park, NC, to review the draft technical report on toxicity studies of glyphosate.

Dr. Po Chan, NIEHS, introduced the short-term toxicity studies of glyphosate by reviewing the uses and rationale for the study, findings from chemical disposition studies, experimental design, and results.

Dr. Garman, a principal reviewer, said that the report was thoroughly prepared and detailed, and that it did an excellent job reviewing the background for the study and the available literature on glyphosate. He added that the isoproterenol/propranolol study included in the report is quite interesting and helps establish the mechanism for salivary gland alteration.

Dr. Garman said that certain details of the salivary gland alteration study should be clarified, namely, which type of glandular acinus within the submandibular salivary gland was most affected by glyphosate, and whether, in Table 11, only the parotid salivary gland was assayed in measuring the severity of changes brought on by glyphosate treatment. Dr. J. Mahler, NIEHS, said the severity grades were based on the parotid glands only.

Dr. Goodman, another principal reviewer, said the report was well-written. He suggested that the the lack of of any reproductive toxicity attributable to glyphosate treatment was an important finding and should be included in the abstract of the report.

After further discussion of editorial matters, Dr. Longnecker accepted the report on behalf of the panel.